

## MENINGOCOCCAL VACCINE WAIVER FORM

The Advisory Committee on Immunization Practices (ACIP) of the U.S. Centers for Disease Control and Prevention (CDC) recommends that first-year college students living in residence halls should receive at least one (1) dose of meningococcal conjugate vaccine before college entry. All students who will be living in campus housing and who are age 23 or younger must submit proof of one (1) dose of meningococcal conjugate vaccine.

Fayetteville State University requires that every student residing in a Housing & Residence Life facility has received a one-time vaccination against meningococcal disease or has signed a waiver. This waiver will detail information on the risks associated with meningococcal disease, the availability, and the effectiveness of vaccine, and that they choose not to be vaccinated for religious or other reasons. Please read below and complete **ONE OF THE TWO** sections below, as it pertains to you.

### MENINGOCOCCAL VACCINE REQUIREMENT

I have received the meningococcal vaccine as required by Fayetteville State University Housing & Residence Life for individuals residing in University Housing. Documentation from a physician or health clinic or receipt of vaccine, and date vaccine was administered is has been upload into SHS Patient Portal.

\_\_\_\_\_  
Signature of Patient or Legal Representative

\_\_\_\_\_  
Date

\_\_\_\_\_  
Bronco ID #

\_\_\_\_\_  
Age

### WAIVER AGE 18 YEARS OR OLDER

I am 18 years of age or older. I have received and reviewed the information provided on the risk of meningococcal disease and the effectiveness and availability of meningococcal vaccine. I understand that meningococcal disease is a rare but life-threatening illness. I understand that Fayetteville State University requires that an individual enrolled in an institute of higher education in North Carolina who resides in University Housing shall receive vaccination against meningococcal disease unless the individual signs the waiver to the vaccination.

I voluntarily agree to release, discharge, indemnify, and hold harmless FSU from any and all costs, liabilities, expenses, claims, demands, or causes of action on account of any loss or personal injury that might result from my non-compliance with the Universities requirements. I choose to waive receipt of the meningococcal vaccine.

#### Risks and Benefits of Meningococcal Vaccination

<i>Vaccine-Preventable Disease</i>	<i>Effectiveness of Vaccine</i>	<i>Possible Side Effects of Vaccination</i>
Meningococcal disease is an acute, potentially severe illness that most often causes meningitis, an infection of the spinal fluid and the fluid that surrounds the brain. It leads to sudden onset of fever, headache, and stiff neck and is usually accompanied by nausea, vomiting, light sensitivity, and altered mental status. Less commonly, it can cause pneumonia, arthritis, and ear/throat infections. Meningococcal disease can result in hearing loss, nervous system problems, seizures, strokes, loss of limbs (arms, legs), and even death.	A protective level of antibody is usually achieved within 7 – 10 days of vaccination. The vaccines protect about 90% of individuals who get them.	The most common side effects are redness or pain at the injection site lasting 1– 2 days, headache, and fatigue. Serious allergic reactions are very rare.

\_\_\_\_\_  
Signature of Patient or Legal Representative

\_\_\_\_\_  
Date

\_\_\_\_\_  
Bronco ID #

\_\_\_\_\_  
Age

# Meningococcal ACWY Vaccine:

## What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

**Meningococcal ACWY vaccine** can help protect against **meningococcal disease** caused by serogroups A, C, W, and Y. A different meningococcal vaccine is available that can help protect against serogroup B.

**Meningococcal disease** can cause meningitis (infection of the lining of the brain and spinal cord) and infections of the blood. Even when it is treated, meningococcal disease kills 10 to 15 infected people out of 100. And of those who survive, about 10 to 20 out of every 100 will suffer disabilities such as hearing loss, brain damage, kidney damage, loss of limbs, nervous system problems, or severe scars from skin grafts.

Meningococcal disease is rare and has declined in the United States since the 1990s. However, it is a severe disease with a significant risk of death or lasting disabilities in people who get it.

Anyone can get meningococcal disease. Certain people are at increased risk, including:

- Infants younger than one year old
- Adolescents and young adults 16 through 23 years old
- People with certain medical conditions that affect the immune system
- Microbiologists who routinely work with isolates of *N. meningitidis*, the bacteria that cause meningococcal disease
- People at risk because of an outbreak in their community

### 2. Meningococcal ACWY vaccine

**Adolescents** need 2 doses of a meningococcal ACWY vaccine:

- First dose: 11 or 12 years of age
- Second (booster) dose: 16 years of age

In addition to routine vaccination for adolescents, meningococcal ACWY vaccine is also recommended for **certain groups of people**:

- People at risk because of a serogroup A, C, W, or Y meningococcal disease outbreak
- People with HIV
- Anyone whose spleen is damaged or has been removed, including people with sickle cell disease
- Anyone with a rare immune system condition called “complement component deficiency”
- Anyone taking a type of drug called a “complement inhibitor,” such as eculizumab (also called “Soliris”®) or ravulizumab (also called “Ultomiris”®)
- Microbiologists who routinely work with isolates of *N. meningitidis*
- Anyone traveling to or living in a part of the world where meningococcal disease is common, such as parts of Africa
- College freshmen living in residence halls who have not been completely vaccinated with meningococcal ACWY vaccine
- U.S. military recruits



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention

---

### 3. Talk with your health care provider

---

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of meningococcal ACWY vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone meningococcal ACWY vaccination until a future visit.

There is limited information on the risks of this vaccine for pregnant or breastfeeding people, but no safety concerns have been identified. A pregnant or breastfeeding person should be vaccinated if indicated.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting meningococcal ACWY vaccine.

Your health care provider can give you more information.

---

### 4. Risks of a vaccine reaction

---

- Redness or soreness where the shot is given can happen after meningococcal ACWY vaccination.
- A small percentage of people who receive meningococcal ACWY vaccine experience muscle pain, headache, or tiredness.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

### 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

### 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

### 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).

